Robust Computational Modeling of Human Ventricular Activation

2017 CASIS Workshop

Jonathan Cranford Postdoctoral Research Staff Member




LLNL-PRES-731201

Lawrence Livermore National Laboratory

A Mechanical and Electrical System

Every life-sustaining heartbeat must have coordinated contraction of the ventricles




"Cardiac-Cycle-Animated" by DrJanaOfficial [CC BY-SA 4.0 (http://creativecommons.org/licenses/by-sa/4.0]], via Wikimedia Commons (modified)



A Mechanical and Electrical System

Every life-sustaining heartbeat must have coordinated contraction of the ventricles




"Latidos" by josiño (Own work) [Public domain], via Wikimedia Commons





A Mechanical and Electrical System

Every life-sustaining heartbeat must have coordinated contraction of the ventricles




"Latidos" by josiño (Own work) [Public domain], via Wikimedia Commons



A Mechanical and Electrical System

Every life-sustaining heartbeat must have coordinated contraction of the ventricles



<u>Abbreviations</u>

RV = right ventricle

LV = left ventricle

S = Septum

W = Wall

RA = right atrium

LA = left atrium

Coordinated ventricular contraction is regulated by electrical depolarization



"ECG Principle fast" by Kalumet (selbst erstellt = Own work) [GFDL (http://www.gnu.org/copyleft/fdl.html) or CC-BY-A-3.0 (http://creativecommons.org/licenses/by-sa/3.0/)], via Wikimedia Commons (modified)


The heart's electrical system drives the mechanical system





A Mechanical and Electrical System

Every life-sustaining heartbeat must have coordinated contraction of the ventricles



<u>Abbreviations</u>

RV = right ventricle

LV = left ventricle


S = Septum

W = Wall


RA = right atrium

LA = left atrium

Coordinated ventricular contraction is regulated by electrical depolarization



Ventricular depolarization cascade is initiated by the Purkinje network




"ECG Principle fast" by Kalumet (selbst erstellt = Own work) [GFDL (http://www.gnu.org/copyleft/fdl.html) or CC-BY-A-3.0 (http://creativecommons.org/licenses/by-sa/3.0/)], via Wikimedia Commons (modified)

"Conductionsystemoftheheart" by Madhero88 (Own work) [CC BY 3.0 (http://creativecommons.org/licenses/by/3.0)], via Wikimedia Commons (modified)

The heart's electrical system drives the mechanical system



The Purkinje Network



"Conductionsystemoftheheart" by Madhero88 (Own work) [CC BY 3.0 (http://creativecommons.org/licenses/by/3.0)], via Wikimedia Commons (modified)


Is an "insulating superhighway" until the very ends

At the ends are the Purkinje-myocyte junctions (PMJs)

PMJs deliver charge to a large amount of tissue nearly synchronously

PMJs provide critical initial conditions for electrical wave propagation

The Purkinje Network




"Conductionsystemoftheheart" by Madhero88 (Own work) [CC BY 3.0 (http://creativecommons.org/licenses/by/3.0)], via Wikimedia Commons (modified)

Is an "insulating superhighway" until the very ends

At the ends are the Purkinje-myocyte junctions (PMJs)

PMJs deliver charge to a large amount of tissue nearly synchronously


PMJs provide critical initial conditions for electrical wave propagation



PMJs provide proper initial conditions critical for physiological electrical wave propagation, and thus, regulate proper contraction



Modeling the Purkinje Network




"Conductionsystemoftheheart" by Madhero88 (Own work) [CC BY 3.0 (http://creativecommons.org/licenses/by/3.0)], via Wikimedia Commons (modified)

Full Purkinje network modeling

Fractal trees

Pros: automate/parameterize/optimize network Cons: irregular geometry breakdown, portability

Modeling the Purkinje Network



"Conductionsystemoftheheart" by Madhero88 (Own work) [CC BY 3.0 (http://creativecommons.org/licenses/by/3.0)], via Wikimedia Commons (modified)

Full Purkinje network modeling

Fractal trees

Pros: automate/parameterize/optimize network Cons: irregular geometry breakdown, portability


<u>Superimpose Purkinje dissection images</u>

Pros: anatomical basis

Cons: anatomically nonspecific (Purkinje images

from other ventricles/animals), time intensive

Modeling the Purkinje Network



Full Purkinje network modeling

Fractal trees

Pros: automate/parameterize/optimize network Cons: irregular geometry breakdown, portability

Superimpose Purkinje dissection images

Pros: anatomical basis

Cons: anatomically nonspecific (Purkinje images

from other ventricles/animals), time intensive

"Conductionsystemoftheheart" by Madhero88 (Own work) [CC BY 3.0 (http://creativecommons.org/licenses/by/3.0)], via Wikimedia Commons (modified)

Why not just model the "business end" of the Purkinje network: the PMJs?

Model PMJs as a collection of current injection "stimuli" with location and timing properties

Pros: easily implementable, portable, anatomical basis (ventricular activation timing experiments)

Cons: limited use in investigating Purkinje network/cell defects

Missing piece: robust stimulus protocol to simulate healthy human heart





Our Model of PMJs

Our location/timing of stimuli based on Durrer et al. experiments¹

- used 870 in-tissue electrodes on 7 isolated human hearts
- measured activation timing throughout ventricles


Identified 4 early activation regions






| Region | S | LASW | LPIW | RAIW |
|-------------|---|------|------|------|
| Timing (ms) | 0 | 0 | 0 | 5 |


¹D. Durrer, R.Th.V. Dam, G.E. Freud, M.J. Janse, F.L. Meijler, R.C. Arzbaecher. (1970) Circulation, 41, 899-912





Sensitivity analysis

How many stimuli need
to accurately approximate
early activation regions?










Sensitivity analysis


How many stimuli need


to accurately approximate


early activation regions?



















Sensitivity analysis


How many stimuli need
to accurately approximate
early activation regions?
























Sensitivity analysis


How many stimuli need
to accurately approximate
early activation regions?



















Sensitivity analysis

How many stimuli need to **accurately approximate** early activation regions?

What are the metrics/criteria?

Metrics/criteria derived from ECG (connection to clinic)




Sensitivity analysis


How many stimuli need to **accurately approximate** early activation regions?

What are the metrics/criteria?


Metrics/criteria derived from ECG (connection to clinic)



Sensitivity analysis



Metrics/criteria derived from ECG (connection to clinic)





Sensitivity analysis

How many stimuli need to **accurately approximate** early activation regions?

What are the metrics/criteria?


Metrics/criteria derived from ECG (connection to clinic)





Modeling Electrophysiology

LLNL's Cardioid code: near perfect scaling to over 1M cores



Geometry from Visible Human Project®


Rxn-diffusion equation

$$C_m \frac{\partial V_m}{\partial t} = \frac{1}{\beta} \nabla \cdot (\mathbf{D} \nabla V_m) - I_{ion} + I_{stim}$$
 Capacitive Diffusion Rxn Early charging activation

stimuli

Modeling Electrophysiology

LLNL's Cardioid code: near perfect scaling to over 1M cores



Rxn-diffusion equation


$$C_m \frac{\partial V_m}{\partial t} = \frac{1}{\beta} \nabla \cdot (\mathbf{D} \nabla V_m) - I_{ion} + I_{stim}$$
Capacitive Diffusion Rxn Early activation

ECG code

"Wraps" torso volume conductor around ventricles


Human Project®

Solves Poisson's equation in torso domain using V_m in ventricles as source



stimuli

Sensitivity Analysis Results



Observations

Use all 4 seed stimuli

→ metrics in
physiological
range

Desensitization to using more and more stimuli


28 or more stimuli


→ Metrics comfortably in physiological range


28 stimuli protocol is reasonable middle ground to reproduce physiological metrics



Sensitivity Analysis Results








Adding 27 stimuli to 1 stimulus → big change in ECG


Adding 356 stimuli to 28 stimuli -> relatively smaller change ECG



28 stimuli protocol is reasonable middle ground to reproduce physiological metrics



Ventricular Activation Timings




28 stimuli protocol qualitatively reproduces early activation regions seen in experiments




Number Versus Coverage of Stimuli

Inter-stimuli distance of 0.3 cm (0.5 cm for septum)




Number Versus Coverage of Stimuli

Inter-stimuli distance of 0.3 cm (0.5 cm for septum)




Inter-stimuli distance of 0.7 cm (1.6 cm for septum)




12 "sparse approximation"


Number Versus Coverage of Stimuli

Inter-stimuli distance of 0.3 cm (0.5 cm for septum)











Inter-stimuli distance of 0.7 cm (1.6 cm for septum)



12 "sparse approximation"



Coverage of stimuli dominates over shear number of stimuli



Conclusions

12-stimulus protocol reproduces salient features of ECG seen in clinic and regions of ventricular early activation seen in experiments

Based on sensitivity, we submit that *any* protocol with 3 stimuli located in each of the 4 key regions of early activation, and inter-stimuli distance of 0.7 cm (1.6 cm for septum) is a reasonable activation protocol and model of PMJs

Do we need to keep modeling denser and denser Purkinje networks? Maybe not

Do we need Purkinje network models at all?

Yes, for studying defects in the network and defects of the Purkinje cells themselves No, use 12-stimulus protocol for studying defects downstream of early activation

Future: test robustness on different heart/torso geometries, and on different diseases

We hope the these simple guidelines for modeling activation allow other researchers to quickly and accurately model ventricular activation, enabling more time to develop patient-specific models of downstream disorders, ultimately to create tailored electrophysiological therapies

Acknowledgements

Thank You

All Individuals in Audience

Acknowledgements

Thank You

All Individuals in Audience

LLNL Coauthors

Robert Blake

Jean-Luc Fattebert

James (Jim) Glosli

Felice Lightstone

Thomas O'Hara

David Richards

Sergio Wong

Xiaohua Zhang

External Coauthors

Omar Hafez (UC Davis)

David Krummen (UC San Diego, VA San Diego

Healthcare System)

Joseph Loscalzo (HMS, Brigham and Women's)

Andrew McCulloch (UC San Diego)

Christopher Villongco (UC San Diego)

Acknowledgements

Thank You

All Individuals in Audience

LLNL Coauthors

Robert Blake

Jean-Luc Fattebert

James (Jim) Glosli

Felice Lightstone

Thomas O'Hara

David Richards

Sergio Wong

Xiaohua Zhang

External Coauthors

Omar Hafez (UC Davis)

David Krummen (UC San Diego, VA San Diego

Healthcare System)

Joseph Loscalzo (HMS, Brigham and Women's)

Andrew McCulloch (UC San Diego)

Christopher Villongco (UC San Diego)

LLNL Administrative and Technical Support

Vanessa Gucho

Kim Hegman

Joanne Strickland

HPC staff

CED staff

BBTD staff

LLNL Mentors and

Technical Advisors

Erik Draeger

Ben Fasenfest

Alan Kaplan

William (Liam) Krauss

Sean Lehman

Funding and Resources

LLNL/NNSA LDRD program

15-SI-002 and 13-ERD-035

LLNL Computing Grand

Challenge Program

